

# Potent Anti-Malaria Salts, Co-Crystals and Derivatives of Aminoquinolines with Hydroxyaromatic Acids

Antimalarial activities of salts of aminoquinolines with hydroxyaromatic acids were found to be higher than the parent components. The IC<sub>50</sub> values of the salts were in the range of 5.9-20.3  $\mu$ M against the chloroquine-resistant strain. Antimalarial actions on the parasites were found to be independent of iron chelation, inhibition of haemozoin formation, as well as independent of the anti-oxidant activity of the salts. A covalently linked amide containing compound N-(quinolin-8-yl)-2-(quinolin-8-yloxy)acetamide was identified to have potency similar to the other salts studied in the article.

## Abbreviations

PXRD: Powder X-ray Diffraction; NMR: Nuclear Magnetic Resonance; FT-IR: Fourier Transform Infra-Red; pKa: Negative logarithm of acid dissociation; S.D.: Standard Deviation;  $\mu$ M=Micro-molar; IC<sub>50</sub>: Drug concentration to cause 50% inhibition of the antimalarial activity; FCR: Fc receptor which is a protein; DPPH: 1,1-diphenyl-2-picrylhydrazyl; ELISA: Enzyme-linked immune sorbent assay; 3D7: *P. falciparum* nuclear genome; RPMI-1640 culture media: Roswell Park Memorial Institute 1640 Medium; CCDC: Cambridge Crystallographic database Cambridge

## Introduction

Several organic compounds showing anti-malarial activities such as aminoquinolines [1], 3,4,5-hydroxybenzoic acid [2], 2-hydroxybenzoic acid derivatives [3] and hydroxynaphthoic acid [4] have clear and distinct acid base properties as well as hydrogen bond abilities. Due to acid-base properties associated with quinolines and carboxylic acids, they are suitable to form salts and co-crystals [5,6]. Formation of a salt between two active pharmaceutical ingredients leaves scope to bring together two potent molecules easily to make a stable formulation which may act altogether in different manners. Recently activity of several antimalarial drug molecules have been reviewed and showed how such combinations in the form of co-crystal formed through weak interactions

can alter antimalarial drug activities [7]. On other hand protein–myosin a complex has important role in regulating depletion of red blood cells caused by the Plasmodium during malaria [8]. Thus, it may be expected that peptide mimics of such protein complex may help to develop in drugs that can disrupt the invasion motor [8]. Hence quinoline molecules with peptide bond may be of interest to test for drug activity towards malaria, as some of them may have additional effect due to the peptide unit attached to the quinoline unit. In this context a quinoline amide derivative was shown to adopt hair-pin structure [9] which is a common feature found in many small oligo-peptide compounds which easily adopt hair-pin structure [10-12]. Based on these points, we have studied anti-malarial activities of a series of salts, co-crystal of aminoquinolines with hydroxyaromatic carboxylic acids and also functionalized quinoline derivatives to distinguish their differences in antimalarial activities. The list of compounds, salts that are studied is listed in Chart 1.

## Materials and Method

### ■ Synthesis and characterization of the salts and co-crystal

Salts I-III and co-crystal IV were prepared by crystallization of solution of 1:1 mole ratio of the corresponding aminoquinoline and respective hydroxyaromatic acid. In each case, aminoquinoline and hydroxyaromatic acid was

Baruah JB<sup>1\*</sup>, Khakhlyar P<sup>1</sup>, Holland S<sup>2</sup> & van Zyl RL<sup>2</sup>

<sup>1</sup>Department of Chemistry, Indian Institute of Technology Guwahati, Guwahati 781 039, Assam, India

<sup>2</sup>Pharmacology Division, Department of Pharmacy and Pharmacology, Faculty of Health Sciences; WITS Research, Institute for Malaria (WRIM), University of the Witwatersrand, South Africa

\*Author for correspondence: iuba@iita.ernet.in

dissolved in methanol and kept undisturbed for crystallization. Spectroscopic and analytical details of the salts and co-crystal are as follows: Salt I: Elemental anal Calcd. For  $C_{16}H_{16}N_2O_4$ , C, 63.99; H, 5.37; N, 9.33; and found C, 63.97; H, 5.41, N 9.46. <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): 8.79 (d, J=4.0 Hz, 1H), 8.59 (d, J=8.4 Hz, 1H), 7.79 (d, J=7.6 Hz, 1H), 7.49-7.37 (m, 3H), 7.41 (q, J=4.0 Hz, 1H), 7.19(d, J=8.4 Hz, 1H), 6.93 (d, J=8.0 Hz, 2H), 6.73(d, J=8.0 Hz, 2H), 4.18 (s, 2H). IR (KBr, cm<sup>-1</sup>): 3444 (s), 3321(m), 3082 (w), 1627 (s), 1589 (s), 1566 (m), 1482 (m), 1454 (m), 1432 (w), 1384 (m), 1361(m), 1330 (m), 1305 (w), 1252 (w), 1138 (w), 1028 (w), 859 (w), 787 (m), 702 (w), 664 (w). Salt II: Elemental anal Calcd. For  $C_{16}H_{16}N_2O_6$ , C, 57.83, H, 4.85; N, 8.43; found C, 57.79, H, 4.88; N, 8.45. <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): 9.23 (s, 1H), 8.76 (d, J=4.0 Hz, 1H), 8.53 (d, J=8.4 Hz, 1H), 7.43 (t, J=8.0 Hz, 2H), 7.36 (q, J=4.0 Hz, 1H), 7.19 (d, J=8.0 Hz, 1H), 6.94 (s, 3H), 6.72 (d, J=7.6 Hz, 2H), 5.9 (s, 2H). IR (KBr, cm<sup>-1</sup>): 3458 (s), 3348 (s), 3241 (m), 3076 (w), 2671 (w), 1644 (w), 1599 (w), 1526 (m), 1471 (w), 1360 (s), 1307 (m), 1228 (m), 1179 (w), 1102 (w), 1035 (m), 886 (w), 786 (m), 746 (w), 703 (w), 575 (w). Salt III: Elemental anal Calcd. for  $C_{20}H_{16}N_2O_3$ , C, 72.28; H, 4.85; N, 8.43; found C, 72.33; H, 4.86; N,8.69. <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>, 400MHz): 8.72 (d, J=8Hz, 1H), 8.30 (d, J=8Hz, 1H), 8.17 (d, J=8.4 Hz, 1H), 7.90 (d, J=8 Hz, 1H), 7.76 (d, J=8.8 Hz, 1H), 7.67 (t, J=1.2, 1H), 7.59 (t, J=8.4 Hz, 1H), 7.46-7.39 (m, 1H), 7.39 (d, J=8.4 Hz, 1H), 7.31 (t, J=7.6 Hz, 1H), 7.07 (d, J=8Hz, 1H), 6.89 (d, J=7.6 Hz, 1 H). IR (KBr, cm<sup>-1</sup>): 3424 (m), 3210 (m), 2923 (w), 1631 (m), 1582 (s), 1504 (m), 1467 (m), 1405 (s), 1377 (m), 1328 (m), 1309 (m), 1213 (w), 1017 (w), 863 (w), 815 (m), 795 (m), 771 (m), 693 (m). Cocrystal IV: Elemental anal Calcd. for  $C_{16}H_{14}N_2O_5$ ; C, 61.14; H, 4.49; N, 8.91; found C, 61.20; H, 4.47; N, 9.10. <sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): 9.23 (s, 1H),8.87 (s, 1H), 8.71 (dd, J=1.6 Hz, J=2.4 Hz,1H), 8.17 (dd, J=1.6 Hz, J=6.8 Hz, 1H), 7.46 (q, J=4.0 Hz 1H), 7.30 (t, J=8.0 Hz, 1H), 7.06 (dd, J=0.8 Hz, 1H), 6.91 (s, 2H), 6.86 (dd, J=1.2 Hz, 1H), 5.90 (s, 1H), 4.09 (s, 3H). IR (KBr, cm<sup>-1</sup>): 3516 (m), 3400 (m), 3309 (m), 1686 (s), 1620 (m), 1519 (w), 1425 (m), 1374 (w), 1324 (s), 1272 (w), 1188 (m), 1032 (m), 900 (w), 857 (w), 825 (w), 787 (w), 766 (w), 718 (w).

## ■ Derivatives of aminoquinolines

Compounds V-VI was prepared by reported procedure [13]. Compound VII was prepared by published procedure [5]. Synthesis of ethyl-4-(2-oxo-2-(quinolin-8-ylamino)ethoxy)benzoate (VIII): To a solution of 8-aminoquinoline (0.72 g, 5 mmol) in dry dichloromethane (20 mL), triethylamine (0.71 mL, 5 mmol) was added. Solution was stirred at 0°C for 15 min, followed by which bromoacetyl bromide (0.52 mL, 6 mmol) was added. Reaction mixture was then stirred overnight and filtered. Filtrate was washed with water (10 mL), dried over anhydrous sodium sulphate and then the solvent was removed under reduced pressure. Product was obtained as a brown solid that was purified by recrystallisation from dichloromethane. Amide obtained from above reaction (0.53 g, 2 mmol), ethyl-4-hydroxybenzoate (1.66 g, 10 mmol) and potassium carbonate (1.07 g, 7 mmol) were added to dry acetone (20 mL) under nitrogen and the reaction mixture was stirred at 60°C for 9 h. After completion of reaction solvent was removed under reduced pressure to obtain a pale yellow solid which was recrystallized from methanol. FT-IR (KBr, cm<sup>-1</sup>): 3430 (m), 3323 (m) 2977 (w), 1701(s), 1691(s), 1607 (s), 1544 (s), 1507 (m), 1485 (m), 1427 (w), 1388 (w), 1367 (w), 1330 (w), 1318 (w), 1301 (w), 1279 (m), 1248(s), 1167 (m), 1107 (m), 1055 (m), 841(w), 823 (w), 791 (m), 764 (w), 755 (w), 686 (w), 638 (w), 567 (w), 500 (w). <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 400 MHz): 10.88 (s, 1H), 8.82 (q, J=1.6 Hz, 1H), 8.78 (t, J=4.4 Hz, 1H), 8.14 (dd, J=1.6 Hz, 1H), 8.05 (dd, J=2.0 Hz, 2H), 7.52 (d, J=4.0 Hz, 2H), 7.44 (q, J=4.0 Hz, 1H), 7.11 (d, J=8.8 Hz, 2H), 4.76 (s, 2H), 4.35 (q, J=7.2 Hz, 2H), 1.37(t, J=7.2 Hz, 3H). ESI mass: 351.0266. Calcd. Mass for [M+1]:  $C_{20}H_{19}N_2O_4$ , 351.1345.

Methyl-3-(2-oxo-2-(quinolin-8-ylamino)ethoxy)benzoate (IX): Similar procedure was followed for the synthesis of IX as for VIII except methyl-3-hydroxybenzoate (0.45 g, 3 mmol) was used instead of ethyl-4-hydroxybenzoate. FT-IR (KBr, cm<sup>-1</sup>): 3410 (m), 3332 (m) 2990 (w), 2926 (w), 1708 (s), 1683 (s), 1585 (w), 1541 (s), 1485 (m), 1458 (m), 1435 (m), 1377 (w), 1372 (w), 1313 (w), 1296 (s), 1283 (m), 1222 (w), 1187 (w), 1103 (m), 1075 (m), 988 (w), 889 (w), 827 (w), 793 (w), 758 (m), 733 (m), 678 (w), 600 (w), 576

(w), 550 (w). <sup>1</sup>H-NMR (CDCl<sub>3</sub>, 400MHz): 10.91(s, 1H), 8.84 (q, J=1.2 Hz, 1H), 8.79 (t, J=4.4 Hz, 1H), 8.14 (dd, J=1.6 Hz, 1H), 7.74 (dd, J =1.2 Hz, 2H), 7.52 (d, J=4.4 Hz, 2H), 7.45 (m, 1H), 7.32 (dd, J=2.4 Hz, 2H), 4.76 (s, 2H), 1.21(s, 3H). ESI mass: 337.0094. Calcd. Mass for [M+1]: C<sub>19</sub>H<sub>17</sub>N<sub>2</sub>O<sub>4</sub>, 337.1188.

### ■ In vitro antimalarial assay

Antimalarial activity of the compounds, against the chloroquine-resistant (FCR-3) and chloroquine-sensitive (3D7) strains of *P. falciparum*, were determined using [3H]-hypoxanthine incorporation assay [14,15]. The parasites were continuously maintained in vitro in supplemented RPMI-1640 culture media at a 5% haematocrit. The culture was incubated at 37°C in 5% CO<sub>2</sub>, 3% O<sub>2</sub>, 92% N<sub>2</sub> and synchronized [16] at the ring stage with 5% D-sorbitol before the final parasitaemia and haematocrit were adjusted to 0.5 and 1%, respectively. This suspension was incubated with the test compounds and controls (final concentration of dimethylsulphoxide <1% did not have an inhibitory effect on the parasite) for 24 h at 37°C. Thereafter, 25 µL [3H]-hypoxanthine (Amersham; 1.85 µCi/well) was added to each well. The microtiter plate was further incubated for 24 h; following which the parasitic DNA was harvested onto glass fibre filter mats. The dry mats were then transferred to sample bags containing scintillation fluid (Wallac) and the β-radioactivity counted on the Wallac 1205 Betaplate scintillation counter. Counts per minute (cpm) were generated and the percentage parasite growth calculated taking the uninfected red blood cells into account. The concentration required to inhibit parasite growth by 50% (IC<sub>50</sub> value) was determined from log sigmoid dose response curves using the GraphPad Prism 5.0 software. Chloroquine and quinine were used as the positive controls. Each experiment was repeated, at least, in triplicate.

### ■ Inhibition of β-haematin formation

To ascertain if the mechanism of action was similar to that of chloroquine, the compounds were incubated with 1 mg/mL haemin in 0.5 M acetate buffer with a final pH of 4.7 [17]. The plates were then incubated for 24 h before the unreacted haemin was removed and the β-haematin crystals were quantified by dissolving them in 2 M sodium hydroxide. The

absorbance at 405 nm was used to calculate the IC<sub>50</sub> value at which β-haematin formation was inhibited using GraphPad Prism 5.0 software. Chloroquine was used as positive control and each experiment was repeated three times.

### ■ Iron chelation assay

The ferrozine-iron chelation assay was used to determine the ferrous ion chelating properties of the derivatives [18]. The derivatives were incubated at room temperature in the dark with 0.48 mM ferrous chloride dissolved in 5% (w/v) ammonium acetate for 5 min before 1.2 mM ferrozine was added and further incubated for 30 min. Thereafter the absorbance was measured at 540 nm in an ELISA plate reader (Labsystems Multiskan RC) with Ascent Software (version 2.4). Percentage inhibition of ferrozine-Fe<sup>2+</sup> complex was calculated with ethylenediamine tetra-acetic acid used as a positive control.

### ■ Free radical scavenging assay

The stable free radical-scavenging activity was determined by the 1,1-diphenyl-2-picrylhydrazyl (DPPH•) assay wherein the bleaching rate of a stable free radical, DPPH• was monitored at 540 nm in the presence and absence of the test compounds and positive control, Trolox™. The percentage free radical scavenging activity was determined from at least three independent experiments.

## Results and Discussion

Aminoquinolines generally interact with aromatic carboxylic acids to form hydrogen bonded co-crystals as illustrated by taking example of 6-aminoquinoline with a carboxylic acid shown in Figure 1. In general salt between an acidic and a basic organic compound is formed when ΔpKa.

[ΔpKa = pKa (base) - pKa (acid)] is greater than 2. When ΔpKa value is less than zero, such pair of compounds exclusively forms co-crystal [19,20]. In our case, the differences in pKa (ΔpKa) of corresponding aminoquinoline counterpart with hydroxyaromatic carboxylic acid counterpart in I-IV are 2.49, 0.96, 1.29 and -0.51 respectively. Accordingly, 5-aminoquinoline with 2-hydroxybenzoic acid formed a salt I; 5-aminoquinoline with 3,4,5-trihydroxybenzoic acid formed salt II; 8-aminoquinoline with 1-hydroxy-2-naphthoic acid formed salt III whereas a co-crystal of

8-aminoquinoline with 3,4,5-trihydroxybenzoic acid, IV was formed from reactions of respective hydroxyaromatic acid with corresponding aminoquinoline. Thus, salts I-III and co-crystal IV were confirmed by determining the respective structure by single crystal X-ray diffraction and further confirmed by powder X-ray diffractions (PXRD) technique. To differentiate between the cocrystal and salts, IR spectra of the OH region (2000  $\text{cm}^{-1}$ -4000  $\text{cm}^{-1}$ ) and carbonyl stretching region (1400 $\text{cm}^{-1}$ -1800 $\text{cm}^{-1}$ ) of salt II and co-crystal IV are compared with the corresponding IR spectra of gallic acid. It is found that the OH stretching region of gallic acid and co-crystal IV are broad, have resemblances (Figure 2a), whereas IR spectra of the salt II shows sharp stretching frequencies at 3458  $\text{cm}^{-1}$ , 3348  $\text{cm}^{-1}$  and 33241  $\text{cm}^{-1}$ ; which differentiated the structural differences due to proton transfer to form salt in the case of II. On the other hand, the carbonyl stretching of gallic acid appears at 1703  $\text{cm}^{-1}$ , which appears at 1644  $\text{cm}^{-1}$  and 1686  $\text{cm}^{-1}$  respectively (Figure 2b) in the salt II and in the co-crystal IV. We have confirmed that no overlapping peaks from the quinoline counterpart in the latter region are present.

Quinoline derivatives V-VI were prepared by reacting corresponding acid chloride with aminoquinoline. The compound VII was prepared by a reported procedure [5], whereas compounds VIII-IX were prepared by reaction sequences illustrated in Scheme 1, by reacting aminoquinoline with bromoacetyl bromide followed by reacting the product of this reaction with phenolic compounds. These compounds were characterized from their  $^1\text{H-NMR}$ , IR and mass spectra. Carbonyl frequencies for amide of compound VIII appeared at 1701  $\text{cm}^{-1}$  and 1691  $\text{cm}^{-1}$  and for the compound IX these signals were observed at 1708 and 1683  $\text{cm}^{-1}$ .

### ■ Anti-malaria and pharmacological studies

Salts I-III and co-crystal IV showed anti-malaria activity with  $\text{IC}_{50}$  values in the 6-20  $\mu\text{M}$  range for the chloroquine-resistant strain, but were four fold less active against the chloroquine-sensitive strain (resistance index: 2.0-5.3) (Table 1). However, the activities were less than the quinine standard for both strains. The  $\text{IC}_{50}$  values displayed by 8-aminoquinoline and 5-aminoquinoline [1] were found to be greater than 10  $\mu\text{M}$ . On the other hand, structurally

related compounds such as 6-(8'-pentadecenyl)-2-hydroxybenzoic acid [3] were reported to have  $\text{IC}_{50}$  value 10.1  $\mu\text{M}$ ; for 3,5-dihydroxy-2-naphthoic acid [4] 25  $\mu\text{M}$  and for gallic acid [2] 71.5  $\mu\text{M}$ . These indicate that the  $\text{IC}_{50}$  values for the salts and co-crystals were much less than the individual component. These results point out the combined effects of the two active components influences their effective antimalarial activities and changes the efficiency of the individual component. Derivatisation of aminoquinoline did not enhance antimalarial activity, but instead displayed minimal antimalarial activity at 50  $\mu\text{M}$  when tested against the chloroquine sensitive strain (Table 1). Aminoquinoline compounds are known to chelate metal ions such as iron, thus associated themselves with potential free radical scavenging activity through complex formation [16]. But such antioxidant properties were not retained when the salts, co-crystal and derivatives that were tested. None among these adducts were able to inhibit  $\beta$ -haematin or haemozoin formation which was found to be the case with 2-amino-8-hydroxyquinoline [18-21]. The result is suggestive of the fact that salt of gallic acid (IV) has the lowest  $\text{IC}_{50}$  value with respect to the other tested compounds listed in Table 1. Use of poly-hydroxyaromatic carboxylic acid namely gallic acid in this salt as one component, enhanced water solubility and maintained a neutral condition by neutralizing quinoline counterpart. In order to gain support towards the importance of use of a salt over a covalent compound, we studied the antimalarial activity of compound V-IX (Table 1).  $\text{IC}_{50}$  value of II<IV, in these cases the gallic acid is a common component. Thus the quinoline counterpart affected the antimalarial activity. The salts and co-crystals given in Table 1, which were tested for antimalarial activity retained good safety profiles [1]. Among individual quinoline derivatives the bis-quinoline derivative VII was a found to be a good chelator for iron, and it displayed antimalarial activity comparable to the salts. The other compounds V, VI, VIII, IX were poor in showing antimalarial activity. It is interesting to note that particular compound VII shows least binding affinity to amino acid such as glycine and methionine but have selectivity in binding hydroxy carboxylic acids such as ascorbic acid and lactic acid [5]. From the structural differences among these compounds the compound VII has a hydroxyquinoline part



which probably is responsible for iron-chelation and thereby causing an enhancement in the antimalarial activities. Thus it may be suggested that there is a clear difference between the mechanistic course of action of antimalarial activities by the adducts and the covalently linked compounds.

## Conclusion

We have shown that salts and cocrystal of quinoline with hydroxyaromatic carboxylic acids enhance antimalarial activities over parent compounds. Enhanced potency could be attributed to the increase in hydrophilicity of these salts and co-crystals. Antimalarial potency of covalently linked amides derivatives of aminoquinolines was not observed but incorporation of a hydroxyquinoline part to such a frame has induced potency as in the case of compound VII. It was also observed that the

antimalarial action of salt II was directed against the intra-erythrocytic parasite is of comparable safety profile to chloroquine and its mechanism of action is independent of iron chelation and inhibition on haemozoin formation.

## Acknowledgement

Authors JBB and PK thanks Ministry of Human Resources and Development (India) for departmental support. This work was partially supported by the Faculty of Health Sciences of the University of the Witwatersrand, South Africa and the Belgian Technical Corporation for student support.

## Supporting Information

Crystallographic information files are deposited to Cambridge Crystallographic database, Cambridge and CCDC numbers for I, III and IV are 842526, 843765 and 870031.

## References

- Egan TJ, Hunter R, Kaschula CH, et al. Structure-function relationships in aminoquinolines: effect of amino and chloro groups on quinoline-hematin complex formation, inhibition of beta-hematin formation, and antiplasmodial activity. *J. Med. Chem.* 43(2), 283-291 (2000).
- Ndjonka D, Bergmann B, Agyare C, et al. In vitro activity of extracts and isolated polyphenols from West African medicinal plants against *Plasmodium falciparum*. *Parasitol. Res.* 111(2), 827-834 (2012).
- Chung IM, Seo SH, Kang EY, Park WH, Moon HI. Anti-malarial activity of 6-(8'-Z-pentadecenyl)-salicylic acid from *Viola websteri* in mice. *Malar. J.* 8, 151 (2009).
- Connors R, Schambach F, Read J, et al. Mapping the binding site for gossypol-like inhibitors of *Plasmodium falciparum* lactate dehydrogenase. *Mol. Biochem. Parasitol.* 142(2), 137-142 (2005).
- Kalita D, Deka H, Samanta SS, Guchait S, Baruah JB (2011) Interactions of amino acids, carboxylic acids, and mineral acids with different quinoline derivatives. *J Mol Struct* 990: 183-196.
- Sarma B, Nath NK, Bhogala BR, Nangia A. Synthon Competition and Co-operation in Molecular Salts of Hydroxybenzoic Acids and Aminopyridines. *Cryst. Growth. Des.* 9(3), 1546-1557 (2009).
- Clements M, le Roex T, Blackie M. Multicomponent Crystal Systems of Known Antimalarial Drug Molecules. *Chem. Med. Chem.* 10(11), 1786-1792 (2015).
- Thomas JC, Green JL, Howson RI, et al. Interaction and dynamics of the *Plasmodium falciparum* MTIP-MyoA complex, a key component of the invasion motor in the malaria parasite. *Mol. BioSyst.* 6(3), 494-498 (2010).
- Kalita D, Sarma R, Baruah JB. Formation of symmetry non-equivalent molecules in urea and carbamate derivatives: the role of anion. *Cryst. Eng. Comm.* 11, 803-810 (2009).
- Rajagopal A, Aravinda S, Raghothama S, Shamala N, Balaram P (2011) Chain length effects on helix-hairpin distribution in short peptides with Aid-DALA and Aib-Aib Segments. *Biopolymers (Peptide Sciences)* 96: 744-756.
- Rajagoapl A, Charles BC, Alexey YK, et al. Enhancing the Magnitude of Antibody Responses through Biomaterial Stereochemistry. *ACS. Biomater. Sci. Eng.* 1(7), 601-609 (2015).
- Krishnayan B, Rajagopal A, Raghothama S, Shamala N, Balaram P.  $\beta$ -Turn Analogues in Model  $\alpha\beta$ -Hybrid Peptides: Structural Characterization of Peptides containing  $\beta$ 2.2Ac6c and  $\beta$ 3.3Ac6c Residues. *Chem. Asian. J.* 7(7), 1671-1678 (2012).
- Khakhlary P, Baruah JB. Polymorphs and salts of 4-nitro-N-(quinolin-8-yl)benzamide. *J. Mol. Struct.* 1078, 188-196 (2014).
- Desjardins RE, Canfield CJ, Haynes DJ, Chulay JD. Quantitative assessment of antimalarial activity in vitro by a semi-automated microdilution technique. *Antimic. Agents. Chemother.* 16(6), 710-718 (1979).
- van Zyl RL, Seatlholo ST, Viljoen A. Pharmacological interactions of essential oil constituents on the in vitro growth of *Plasmodium falciparum*. *S. Afr. J. Bot.* 76(4), 662-667 (2010).
- Chemaly SM, Chen CT, van Zyl RL. Naturally occurring cobalamins have antimalarial activity. *J. Inorg. Biochem.* 101(5): 764-773 (2007).
- Gulcin I. Comparison of in vitro antioxidant and antiradical activities of L-tyrosine and L-Dopa. *Amino. Acids.* 32(3), 431-438 (2007).
- van Vuuren JN, van Zyl RL. Abstracts of the 17th World Congress of Basic and Clinical Pharmacology. *Basic. Clin. Pharmacol. Toxicol.* 115(S1), 1-374 (2014).
- Childs SL, Stahly GP, Park A. The salt-cocrystal continuum: the influence of crystal structure on ionization state. *Mol. Pharmaceutics.* 4(3), 323-328 (2007).
- Bhogala BR, Basavoju S, Nangia A. Tape and layer structures in co-crystals of some di- and tricarboxylic acids with 4,4'-bipyridines and isonicotinamide. From binary to ternary co-crystals. *Cryst. Eng. Comm.* 7, 551-562 (2005).
- Hayat F, Moseley E, Salahuddin A, van Zyl RL, Azam A. Antiprotozoal activity of chloroquinoline based chalcones. *Eur. J. Med. Chem.* 46(5), 1897-1905 (2011).